

ANALOGUES OF THE EXPONENTIAL MAP ASSOCIATED WITH COMPLEX STRUCTURES ON NONCOMMUTATIVE TWO-TORI

A. POLISHCHUK

ABSTRACT. We define and study analogues of exponentials for functions on noncommutative two-tori that depend on a choice of a complex structure. The major difference with the commutative case is that our noncommutative exponentials can be defined only for sufficiently small functions. We show that this phenomenon is related to the existence of certain discriminant hypersurfaces in an irrational rotation algebra. As an application of our methods we give a very explicit characterization of connected components in the group of invertible elements of this algebra.

INTRODUCTION

In this paper we study some natural constructions for functions on noncommutative two-tori equipped with a complex structure. Recall that for every number $\theta \in \mathbb{R} \setminus \mathbb{Q}$ the algebra A_θ of smooth functions on the noncommutative torus T_θ (also known as *irrational rotation algebra*) consists of expressions $\sum_{(m,n) \in \mathbb{Z}^2} a_{m,n} U_1^m U_2^n$ where coefficients $a_{m,n} \in \mathbb{C}$ decrease rapidly at infinity and the multiplication is performed using the rule

$$U_1 U_2 = \exp(2\pi i \theta) U_2 U_1.$$

Given an element $\tau \in \mathbb{C} \setminus \mathbb{R}$ (following [6] we will always assume that $\text{Im}(\tau) < 0$) we define a derivation

$$\delta_\tau : A_\theta \rightarrow A_\theta : \sum a_{m,n} U_1^m U_2^n \mapsto 2\pi i \cdot \sum_{m,n} (m\tau + n) a_{m,n} U_1^m U_2^n$$

We consider δ_τ as a complex structure on T_θ and denote the resulting complex noncommutative torus by $T_{\theta,\tau}$.

The main object of our study is the equation

$$\delta_\tau(x) = xa \tag{0.1}$$

for $x \in A_\theta$, where $a \in A_\theta$ is given. In the commutative case this equation is clearly related with the exponential map on smooth functions. It turns out that there is a local analogue of this map for A_θ . However, in noncommutative case there seem to be serious reasons why the exponential map does not extend to all functions. For example, we show that (0.1) has a nonzero solution iff $\text{tr}(a) \in 2\pi i(\mathbb{Z} + \mathbb{Z}\tau)$, where $\text{tr}(\sum a_{m,n} U_1^m U_2^n) = a_{0,0}$ (see Corollary 2.7), but these solutions are not necessarily invertible in A_θ (as in the commutative case).

The study of equation (0.1) turns out to be closely related to the study of holomorphic structures on the trivial holomorphic bundle over $T_{\theta,\tau}$. By a holomorphic bundle on $T_{\theta,\tau}$ we mean a right projective module E over A_θ equipped with a δ_τ -connection, i.e., a linear

Supported in part by NSF grant.

map $\overline{\nabla} : E \rightarrow E$ satisfying the Leibnitz rule $\overline{\nabla}(ea) = \overline{\nabla}(e)a + e\delta_\tau(e)$. The category $\mathcal{C}$ of holomorphic bundles on $T_{\theta,\tau}$ was studied in [6] and [4]. The main result of [4] identifies $\mathcal{C}$ with the heart of a certain t -structure on the derived category of coherent sheaves on the elliptic curve $\mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$. This leads to a classification of holomorphic bundles on $T_{\theta,\tau}$ up to isomorphism. We apply this classification to the study of solutions of (0.1), in particular, to the question of existence of a solution $x \in A_\theta^*$, where $A_\theta^* \subset A_\theta$ is the set of invertible elements. The main result here is Theorem 2.6 stating that the map

$$A_\theta^*/\mathbb{C}^* \rightarrow A_\theta : x \mapsto x^{-1}\delta_\tau(x)$$

identifies $A_\theta^*/\mathbb{C}^*$ with $\Omega_\tau + 2\pi i(\mathbb{Z} + \mathbb{Z}\tau)$ where Ω_τ is a dense open subset in the hyperplane $H = \{a \in A_\theta : \text{tr}(a) = 0\}$. More precisely, we prove that Ω_τ is the complement to the *discriminant hypersurface* $\Theta_\tau \subset H$ consisting of $a \in H$ such that the equation

$$\delta_\tau(x) + ax - xa = 0$$

has a nontrivial solution $x \in H$. We call it a hypersurface since it coincides with the zero locus of a global holomorphic section of a holomorphic line bundle over A_θ (induced by Quillen's determinant line bundle on the space of Fredholm operators of index zero). We study the structure of Θ_τ in more detail. In particular, we present some results that make us believe that Θ_τ should be irreducible in an appropriate sense.

As a byproduct of our study we deduce the following statement about the structure of the group A_θ^* . It is well known that the group of connected components of A_θ^* can be identified with $\mathbb{Z}^2$. We prove that for every $x \in A_\theta^*$ one has $\text{tr}(x^{-1}\delta_\tau(x)) \in 2\pi i(\mathbb{Z} + \mathbb{Z}\tau) \subset \mathbb{C}$ and that the map $x \mapsto \text{tr}(x^{-1}\delta_\tau(x))$ induces an isomorphism on the groups of connected components.

The paper is organized as follows. In section 1 we study formal exponential maps for A_θ . First, we present the construction of a map

$$\mathbf{E}_l(\tau, \cdot) : A_\theta[[t]]/\mathbb{C}[[t]] \rightarrow 1 + tA_\theta[[t]]$$

that specializes to $a \mapsto \exp(ta)/\text{tr}(\exp(ta))$ in the commutative case. Then in section 1.4 we construct and study the normalized map

$$\text{Exp}_l(\tau, \cdot) : A_\theta[[t]] \rightarrow 1 + tA_\theta[[t]]$$

that specializes to $\exp(ta)$ in the commutative case. In section 2 we prove that our formal exponentials converge in a neighborhood of zero and then study logarithmic derivatives of invertible elements and the discriminant hypersurface.

Acknowledgment. I am grateful to Chris Phillips for many useful conversations and to Hanfeng Li for correcting a mistake in the first draft.

1. FORMAL EXPONENTIALS

1.1. Notation. For every $v = (m, n) \in \mathbb{Z}^2$ we set

$$U_v = \exp(-\pi i \theta mn) U_1^m U_2^n$$

We have the following product rule in A_θ :

$$U_v \cdot U_{v'} = \exp(2\pi i \langle v, v' \rangle) U_{v+v'},$$

where for $v' = (m', n')$ we set

$$\langle v, v' \rangle = \langle v, v' \rangle_\theta = \frac{1}{2}\theta(mn' - m'n).$$

For an element $a = \sum_{v \in \mathbb{Z}^2} a_v U_v \in A_\theta$ we define the *support* of a as the set of all $v \in \mathbb{Z}^2$ such that $a_v \neq 0$. Note that elements supported on a fixed rank-1 subgroup $\mathbb{Z} \subset \mathbb{Z}^2$ form a commutative subalgebra in A_θ .

Recall that A_θ is equipped with the $\mathbb{C}$ -antilinear antiinvolution $*$ defined by $(U_1)^* = U_1^{-1}$, $(U_2)^* = U_2^{-1}$. For $v \in \mathbb{Z}^2$ we have $(U_v)^* = U_{-v}$, hence for $a = \sum_{v \in \mathbb{Z}^2} a_v U_v$ we get $a^* = \sum_v a_v^* U_v$, where $a_v^* = \overline{a_{-v}}$.

Let $\iota : \mathbb{Z}^2 \rightarrow \mathbb{C}$ be the homomorphism sending (m, n) to $2\pi i(m\tau + n)$. Then for every $v \in \mathbb{Z}^2$ we have

$$\delta_\tau(U_v) = \iota(v) \cdot U_v.$$

1.2. First construction. Our formal exponentials will live in the ring $A_\theta[[t]]$, where t is a formal variable commuting with A_θ . We extend δ_τ to a $\mathbb{C}[[t]]$ -linear derivation of $A_\theta[[t]]$ and the trace to a $\mathbb{C}[[t]]$ -linear functional $\text{tr} : A_\theta[[t]] \rightarrow \mathbb{C}[[t]]$.

Theorem 1.1. *For every $\tau \in \mathbb{C} \setminus \mathbb{R}$ there exists a unique map*

$$\mathbf{E}_l(\tau, \cdot) : A_\theta[[t]]/\mathbb{C}[[t]] \rightarrow 1 + tA_\theta[[t]],$$

with the following properties:

- (i) $\delta_\tau(\mathbf{E}_l(\tau, a)) = t\mathbf{E}_l(\tau, a) \cdot \delta_\tau(a)$;
- (ii) $\text{tr}(\mathbf{E}_l(\tau, a)) = 1$.

We need the following purely algebraic statement.

Lemma 1.2. *Let A be an associative algebra over $\mathbb{Q}$, M an A -bimodule, and $d : A \rightarrow M$ a derivation. Assume that $a_1, \dots, a_n$ are elements of A such that*

$$d(a_i) = a_{i-1}d(a_1) \text{ for } i = 2, \dots, n.$$

Then

$$a_n d(a_1) \in [A, M] + d(A),$$

where $[A, M] \subset M$ is the linear span of the elements of the form $[a, m] = am - ma$ for $a \in A, m \in M$.

Proof. For every $n_1, \dots, n_k \in [1, n]$ let us denote

$$[n_1, \dots, n_k] = a_{n_1} \dots a_{n_{k-1}} d(a_{n_k}) \bmod [A, M] + d(A)$$

From the Leibnitz identity we get

$$[n_1, \dots, n_k] + c.p.(1, \dots, k) = 0$$

in $A/([A, M] + d(A))$, where $c.p.(1, \dots, k)$ denotes the terms obtained by cyclic permutation from $(1, \dots, k)$. Since k is invertible in A , this implies that

$$\sum_{n_1 + \dots + n_k = n} [n_1, \dots, n_k] = 0. \tag{1.1}$$

On the other hand, the identity $d(a_i) = a_{i-1}d(a_1)$ implies that

$$[n_1, \dots, n_k] = [n_1, \dots, n_{k-1}, n_k - 1, 1] \quad (1.2)$$

for $n_k > 1$. Let us denote

$$b_k = \sum_{n_1 + \dots + n_k = n} [n_1, \dots, n_k, 1].$$

For example, $b_1 = [n, 1] = a_n d(a_1)$, $b_{n+1} = 0$. We claim that for every k

$$b_k = -b_{k-1}.$$

Indeed, using (1.2) we get

$$b_k = \sum_{n_1 + \dots + n_k = n} [n_1, \dots, n_{k-1}, n_k + 1] = \sum_{n_1 + \dots + n_k = n+1, n_k > 1} [n_1, \dots, n_k].$$

Now applying (1.1) we can replace this with

$$b_k = - \sum_{n_1 + \dots + n_k = n+1, n_k = 1} [n_1, \dots, n_k] = -b_{k-1}$$

which proves our claim. Therefore,

$$a_n d(a_1) = b_1 = \pm b_{n+1} = 0$$

modulo $[A, M] + d(M)$. □

Proof of Theorem 1.1. Let us first show the uniqueness of $\mathbf{E}_l(\tau, a)$. If $x, y \in 1 + tA_\theta[[t]]$ satisfy $x^{-1}\delta(x) = y^{-1}\delta(y) = t\delta(a)$ then

$$\delta(xy^{-1}) = \delta(x)y^{-1} - xy^{-1}\delta(y)y^{-1} = 0$$

hence $x = cy$ with $c \in \mathbb{C}[[t]]$. Furthermore,

$$1 = \text{tr}(x) = \text{tr}(cy) = c \text{tr}(y) = c,$$

so $x = y$. To show the existence let us write $\mathbf{E}_l(\tau, a) = 1 + a_1 t + a_2 t^2 + \dots$ with some $a_i \in A_\theta[[t]]$, such that $\text{tr}(a_i) = 0$ (so that condition (ii) is satisfied) and such that

$$\delta_\tau(a_i) = a_{i-1} \delta_\tau(a) \text{ for } i \geq 1,$$

where $a_0 = 1$. It is clear that condition (i) would follow, so it is enough to show the existence of such a_i 's. To this end we set $a_1 = a - \text{tr}(a)$ and apply Lemma 1.2 to construct a_i for $i \geq 2$ inductively. Note that we have

$$[A_\theta, A_\theta] \subset H = \delta_\tau(A_\theta)$$

where $H \subset A_\theta$ is the space of elements x with $\text{tr}(x) = 0$. A similar inclusion holds for the ring of formal power series. Thus, when $a_1, \dots, a_n$ are already constructed, Lemma 1.2 allows us to conclude that $a_n \delta_\tau(a) \in \delta_\tau(A_\theta[[t]])$, so we can define a_{n+1} as the unique element satisfying $\delta_\tau(a_{n+1}) = a_n \delta_\tau(a)$ and $\text{tr}(a_{n+1}) = 0$. □

Remark. In the commutative case we have

$$\mathbf{E}_l(\tau, a) = \exp(ta) / \text{tr}(\exp(ta))$$

due to property (ii). In section 1.4 we will present a way of normalizing noncommutative exponentials which reduces to the standard normalization in the commutative case.

Similarly, one can define the function $\mathbf{E}_r(\tau, \cdot)$ such that

$$\delta_\tau(\mathbf{E}_r(\tau, a)) = t\delta_\tau(a) \cdot \mathbf{E}_r(\tau, a)$$

and $\text{tr}(\mathbf{E}_r(\tau, a)) = 1$. Let us extend the standard antiinvolution $*$: $A_\theta \rightarrow A_\theta$ to an antiinvolution of $A_\theta[[t]]$ by t -linearity. From the equation $\delta_\tau(a)^* = \delta_{\bar{\tau}}(a^*)$ one can easily derive that

$$\mathbf{E}_r(\tau, a)^* = \mathbf{E}_l(\bar{\tau}, a^*). \quad (1.3)$$

Proposition 1.3. *For every $a \in A_\theta[[t]]$ one has*

$$\mathbf{E}_l(\tau, a) \cdot \mathbf{E}_r(\tau, -a) = \mathbf{E}_r(\tau, -a) \cdot \mathbf{E}_l(\tau, a) =: \mathbf{s}(\tau, a) \in 1 + t\mathbb{C}[[t]].$$

Proof. Indeed, let $x = \mathbf{E}_l(\tau, a)$, $y = \mathbf{E}_r(\tau, -a)$, $\delta = \delta_\tau$. Then

$$\delta(xy) = \delta(x)y + x\delta(y) = tx\delta(a)y - tx\delta(a)y = 0.$$

This implies that $z = xy \in 1 + t\mathbb{C}[[t]]$. Since x is invertible in $A_\theta[[t]]$ and z commutes with x^{-1} , we derive that $y = zx^{-1}$ and so $yx = z = xy$. $\square$

Corollary 1.4. *One has $\overline{\mathbf{s}(\tau, a)} = \mathbf{s}(\bar{\tau}, -a^*)$.*

Proof. We have

$$\overline{\mathbf{s}(\tau, a)} = \overline{\mathbf{E}_r(\tau, -a)^* \mathbf{E}_l(\tau, a)^*} = \mathbf{E}_l(\bar{\tau}, -a^*) \mathbf{E}_r(\bar{\tau}, a^*) = \mathbf{s}(\bar{\tau}, -a^*). \quad \square$$

1.3. Formula in terms of coefficients. Below we are going to write an explicit formula for $\mathbf{E}_l(\tau, a)$ in terms of the coefficients of a .

Recall that $\iota : \mathbb{Z}^2 \rightarrow \mathbb{C}$ is the homomorphism sending (m, n) to $2\pi i(m\tau + n)$. Let us define a collection of symmetric functions $f_n(v_1, \dots, v_n)$ of n lattice vectors recursively by setting $f_0 = f_1(v) \equiv 1$,

$$f_n(v_1, \dots, v_n) = \iota(v_1 + \dots + v_n)^{-1} \sum_{i=1}^n \iota(v_i) f_{n-1}(v_1, \dots, \widehat{v}_i, \dots, v_n) \exp(2\pi i \langle \sum_{j \neq i} v_j, v_i \rangle)$$

if $v_1 + \dots + v_n \neq 0$, and $f_n(v_1, \dots, v_n) = 0$ otherwise. It is easy to see that f_n can also be given by the following formula:

$$f_n(v_1, \dots, v_n) = \sum_{\sigma \in S(v_1, \dots, v_n)} \frac{\iota(v_{\sigma(1)}) \dots \iota(v_{\sigma(n)}) \exp(2\pi i \sum_{i < j} \langle v_{\sigma(i)}, v_{\sigma(j)} \rangle)}{\iota(v_{\sigma(1)}) \iota(v_{\sigma(1)} + v_{\sigma(2)}) \dots \iota(v_{\sigma(1)} + \dots + v_{\sigma(n)})},$$

where the summation is taken over the set $S(v_1, \dots, v_n) \subset S_n$ of all permutations σ for which the denominator in the corresponding term does not vanish. When we want to stress the dependence of f_n on θ we write $f_n(v_1, \dots, v_n; \theta)$ instead of $f_n(v_1, \dots, v_n)$.

Let $\text{Div}(\mathbb{Z}^2)_{\geq 0}$ denote the semigroup of effective divisors on $\mathbb{Z}^2$, i.e., of formal linear combinations $D = n_1(v_1) + \dots + n_k(v_k)$ with $n_i \geq 0$ and $v_i \in \mathbb{Z}^2$. We set

$$\text{supp}(D) = \{v_i : n_i > 0\}$$

$$\begin{aligned}
\deg(D) &= n_1 + \dots + n_k, \\
s(D) &= n_1 v_1 + \dots + n_k v_k \in \mathbb{Z}^2, \\
D! &= n_1! \dots n_k!
\end{aligned}$$

Theorem 1.5. *For an element $a = \sum_{v \neq 0} a_v U_v \in A_\theta[[t]]$, such that $a_0 = 0$, one has*

$$\mathbf{E}_l(\tau, a) = \sum_{D \in \text{Div}(\mathbb{Z}^2)_{\geq 0}} \frac{t^{\deg(D)}}{D!} c(D) a_D U_{s(D)},$$

where for $D = (v_1) + \dots + (v_d)$ (with not necessarily distinct v_i) we set

$$c(D) = c_\theta(D) := f_d(v_1, \dots, v_d; \theta),$$

$$a_D = a_{v_1} \dots a_{v_d}.$$

Proof. Recall that $\mathbf{E}_l(\tau, a) = \sum a_i t^i$, where $\delta(a_i) = a_{i-1} \delta(a)$ and $\text{tr}(a_i) = 0$ for $i \geq 1$. This easily implies that

$$a_n = \sum_{D \in \text{Div}(\mathbb{Z}^2)_{\geq 0} : \deg D = n} \frac{c(D)}{D!} a_D U_{s(D)}$$

for some constants $c(D)$ that do not depend on a . Note that the $c(D)$ are uniquely determined from this equation since the functions $a \mapsto a_D$ for $D \in \text{Div}(\mathbb{Z}^2)_{\geq 0}$ are linearly independent. Note also that we have $c(D) = 0$ if $s(D) = 0$. Comparing the coefficients with $a_D U_{s(D)}$ in the equation $\delta(a_d) = a_{d-1} \delta(a)$, where $d = \deg D$, we obtain the equation

$$\iota(s(D)) \frac{c(D)}{D!} = \sum_{v \in \text{supp}(D)} \iota(v) \frac{c(D-v)}{(D-v)!} \exp(2\pi i \langle s(D-v), v \rangle).$$

Equivalently,

$$\iota(s(D)) c(D) = \sum_{v \in \text{supp}(D)} n_v \iota(v) c(D-v) \exp(2\pi i \langle s(D-v), v \rangle), \quad (1.4)$$

where n_v is the multiplicity of v in D . This implies that $c(D) = f_d(v_1, \dots, v_d)$ for $D = (v_1) + \dots + (v_d)$. $\square$

Corollary 1.6. *For every $v_1, \dots, v_n \in \mathbb{Z}^2$ such that $v_1 + \dots + v_n = 0$ one has*

$$\sum_{i=1}^n \iota(v_i) f_{n-1}(v_1, \dots, \widehat{v_i}, \dots, v_n) = 0.$$

Proof. Apply equation (1.4) to $D = (v_1) + \dots + (v_n)$. $\square$

Remark. We do not know a direct combinatorial proof of the identity of Corollary 1.6 except in the case when $v_1, \dots, v_n$ are sufficiently generic. See the remark after Theorem 1.10.

Recall that $(\sum_v a_v U_v)^* = \sum_v \overline{a_{-v}} U_v$. Therefore, using (1.3) and Theorem 1.5 we get

$$\mathbf{E}_r(\tau, a) = \sum_{D \in \text{Div}(\mathbb{Z}^2)_{\geq 0}} \frac{t^{\deg(D)}}{D!} c_{-\theta}(D) a_D U_{s(D)}.$$

1.4. Normalized exponential. In this section we define a natural modification of $\mathbf{E}_l(\tau, \cdot)$ that reduces to the usual exponential in the commutative case.

The idea is to modify the symmetric functions $f_n(v_1, \dots, v_n)$ that appear in the explicit formula for $\mathbf{E}_l(\tau, \cdot)$. Let us consider the rational functions

$$R_n(x_1, \dots, x_n) = \frac{x_1 x_2 \dots x_n}{x_1(x_1 + x_2) \dots (x_1 + \dots + x_n)}.$$

For a permutation $\sigma \in S_n$ and an n -tuple $x = (x_1, \dots, x_n)$ let us denote $x^\sigma = (x_{\sigma(1)}, \dots, x_{\sigma(n)})$. Recall that for an n -tuple of lattice vectors $\bar{v} = (v_1, \dots, v_n)$ we have

$$f_n(\bar{v}) = \sum_{\sigma \in S(\bar{v})} R_n(\iota(\bar{v})^\sigma) \exp(2\pi i \sum_{i < j} \langle v_{\sigma(i)}, v_{\sigma(j)} \rangle)$$

where $\iota(\bar{v}) = (\iota(v_1), \dots, \iota(v_n))$, and $S(\bar{v}) \subset S_n$ is the set of all permutations σ such that R_n is regular at $\bar{v}^\sigma$. We want to define a modified expression $f_n^*(\bar{v})$ that takes into account contributions from σ such that R_n is not defined at $\bar{v}^\sigma$. For this we will combine several terms corresponding to such permutations, so that the poles will cancel out.

An n -tuple $\bar{v} = (v_1, \dots, v_n)$ determines a map

$$t_{\bar{v}} : S_n \rightarrow \mathbb{Z}^2 \otimes_{\mathbb{Z}} \mathbb{Z}^2 : \sigma \mapsto \sum_{i < j} v_{\sigma(i)} \otimes v_{\sigma(j)}.$$

The fibers of $t_{\bar{v}}$ define a partition

$$S_n = S(\bar{v}, 1) \sqcup \dots \sqcup S(\bar{v}, r)$$

of S_n . The crucial observation is the following remarkable cancellation of poles.

Proposition 1.7. *The rational functions*

$$R_{S(\bar{v}, j)}(x) := \sum_{\sigma \in S(\bar{v}, j)} R_n(x^\sigma)$$

for $j = 1, \dots, r$ are regular at $\iota(\bar{v})$. More precisely, for every collection $1 \leq i_1 < \dots < i_r \leq n$ such that $v_{i_1} + \dots + v_{i_r} = 0$ the functions $R_{S(\bar{v}, j)}$ do not have poles along the hyperplane $x_{i_1} + \dots + x_{i_r} = 0$.

The proof is based on two lemmata.

Lemma 1.8. *If $v_1 + \dots + v_n = 0$ then*

$$\sum_{i < j} v_i \otimes v_j = \sum_{i < j} v_{i+1} \otimes v_{j+1},$$

where we view indices as elements of $\mathbb{Z}/n\mathbb{Z}$, so that $v_{n+1} = v_1$.

The proof is straightforward.

Lemma 1.9. *Let $\zeta \in S_n$ be the cyclic permutation of order n . Then the function*

$$\sum_{i=0}^{n-1} R_n(x^{\zeta^i})$$

has no pole along $x_1 + \dots + x_n = 0$.

Proof. As before we view indices as elements of $\mathbb{Z}/n\mathbb{Z}$. It suffices to prove that the function

$$\sum_{i=1}^n \frac{1}{x_i(x_i + x_{i+1}) \dots (x_i + x_{i+1} + \dots + x_{i+n-1})}$$

has no pole along $x_1 + \dots + x_n = 0$. Since

$$x_i + x_{i+1} + \dots + x_j \equiv -(x_{j+1} + x_{j+2} + \dots + x_{i-1}) \pmod{x_1 + \dots + x_n}$$

we can replace the above expression with

$$\frac{1}{x_1 + \dots + x_n} \cdot \sum_{i=1}^n \frac{(-1)^{i-1}}{(x_1 + \dots + x_{i-1})(x_2 + \dots + x_{i-1})x_{i-1}x_i(x_i + x_{i+1}) \dots (x_i + \dots + x_{n-1})}.$$

Note that the new sum depends only on $(x_1, \dots, x_{n-1})$. Hence, our assertion is equivalent to the identity

$$\sum_{i=1}^n \frac{(-1)^{i-1}}{(x_1 + \dots + x_{i-1})(x_2 + \dots + x_{i-1})x_{i-1} \cdot x_i(x_i + x_{i+1}) \dots (x_i + \dots + x_{n-1})} = 0.$$

Since the left-hand side is homogeneous of degree $-(n-1)$, it suffices to prove that it has no poles. The only possible poles are along the hyperplanes $h_{ij} = x_i + x_{i+1} + \dots + x_j = 0$. Now one can check easily that for every $i \leq j$ there are exactly two terms in the above sum that have poles along $h_{ij} = 0$ and that their polar parts cancel out. $\square$

Proof of Proposition 1.7. We use induction in n and the following recursive formula for R_n :

$$R_n(x_1, \dots, x_n) = R_{n-1}(x_1, \dots, x_{n-1}) \frac{x_n}{x_1 + \dots + x_n}.$$

Assume that the assertion holds for $n-1$. We can write $R_{S(\bar{v}, j)}$ in the form

$$R_{S(\bar{v}, j)} = \sum_{m=1}^n \sum_{\sigma \in S(\bar{v}, j): \sigma(n)=m} R_{n-1}(x_{\sigma(1)}, \dots, x_{\sigma(n-1)}) \frac{x_m}{x_1 + \dots + x_n}$$

For a given m let us denote $S(\bar{v}, j; m)$ the set of $\sigma \in S(\bar{v}, j)$ such that $\sigma(n) = m$. Fix $\sigma_0 \in S(\bar{v}, j; m)$. Then $S(\bar{v}, j; m)\sigma_0^{-1}$ leaves m stable. Furthermore, it is clear that $S(\bar{v}, j; m)\sigma_0^{-1}$ coincides with one piece in the partition of the set of permutations of $(1, \dots, \widehat{m}, \dots, n)$ associated with the $(n-1)$ -tuple $(v_1, \dots, \widehat{v_m}, \dots, v_n)$. Thus, by the induction assumption all the sums

$$\sum_{\sigma \in S(\bar{v}, j; m)} R_{n-1}(x_{\sigma(1)}, \dots, x_{\sigma(n-1)})$$

have no poles along the required hyperplanes. If $v_1 + \dots + v_n \neq 0$ this finishes the proof. It remains to consider the case when $v_1 + \dots + v_n = 0$. By Lemma 1.8 in this case all the sets $S(\bar{v}, i)$ are right cosets for the subgroup $\mathbb{Z}/n\mathbb{Z} \subset S_n$ generated by the cyclic permutation

$\zeta \in S_n$, where $\zeta(i) = i + 1$. Therefore, by Lemma 1.9 the functions $R_{S(\bar{v}, j)}$ have no poles along the hyperplane $x_1 + \dots + x_n = 0$. $\square$

Let us define a rational function of $x = (x_1, \dots, x_n)$ by

$$R_{\bar{v}}(x) = \sum_{\sigma \in S_n} R(x^\sigma) \exp(2\pi i \langle t_{\bar{v}}(\sigma) \rangle) = \sum_{j=1}^r R_{S(\bar{v}, j)}(x) \exp(2\pi i \langle t_{\bar{v}}(S(\bar{v}, j)) \rangle),$$

where $\langle t_{\bar{v}}(S(\bar{v}, j)) \rangle$ is the common value of

$$\langle t_{\bar{v}}(\cdot) \rangle := \langle \cdot, \cdot \rangle \circ t_{\bar{v}} : S_n \rightarrow \mathbb{R}$$

on $S(\bar{v}, j)$. By Proposition 1.7 this function is regular at $\iota(\bar{v})$, so we can set

$$f_n^*(\bar{v}) = f_n^*(\bar{v}; \theta) = R_{\bar{v}}(\iota(\bar{v})).$$

It is easy to check that f_n^* is symmetric in $v_1, \dots, v_n$. As before we have $f_0^* = f_1^*(v) \equiv 1$.

Theorem 1.10. *For an element $a = \sum_v a_v U_v \in A_\theta[[t]]$ consider the series*

$$\text{Exp}_l(\tau, a) := \sum_{D \in \text{Div}(\mathbb{Z}^2)_{\geq 0}} \frac{t^{\deg(D)}}{D!} c^*(D) a_D U_{s(D)} \in 1 + t A_\theta[[t]],$$

where for $D = (v_1) + \dots + (v_d)$ we set

$$c^*(D) = c_\theta^*(D) = f_n^*(v_1, \dots, v_d; \theta).$$

Then

- (i) $\delta_\tau(\text{Exp}_l(\tau, a)) = t \text{Exp}_l(\tau, a) \cdot \delta_\tau(a)$;
- (ii) if $\theta = 0$ or if the support of a is contained in a rank-1 subgroup $\mathbb{Z} \subset \mathbb{Z}^2$ then $\text{Exp}_l(\tau, a) = \exp(ta)$;
- (iii) for $a \in A_\theta[[t]]$ and $z \in \mathbb{C}[[t]]$ we have

$$\text{Exp}_l(z + a) = \exp(tz) \cdot \text{Exp}_l(a). \quad (1.5)$$

Proof. As in Theorem 1.5 the proof of (i) reduces to the following identity:

$$\iota(v_1 + \dots + v_n) f_n^*(v_1, \dots, v_n) = \sum_{m=1}^n \iota(v_m) f_{n-1}^*(v_1, \dots, \widehat{v_m}, \dots, v_n) \exp(2\pi i \langle \sum_{j \neq m} v_j, v_m \rangle).$$

We claim that in fact, there is an identity between the corresponding rational functions:

$$(x_1 + \dots + x_n) R_{\bar{v}}(x_1, \dots, x_n) = \sum_{m=1}^n x_m R_{\bar{v}^{(m)}}(x_1, \dots, \widehat{x_m}, \dots, x_n) \exp(2\pi i \langle \sum_{j \neq m} v_j, v_m \rangle),$$

where $\bar{v}^{(m)} = (v_1, \dots, \widehat{v_m}, \dots, v_n)$. Indeed, we can rewrite the left-hand side as

$$\begin{aligned} (x_1 + \dots + x_n) \sum_{\sigma \in S_n} R_n(x^\sigma) \exp(2\pi i \langle t_{\bar{v}}(\sigma) \rangle) &= \\ \sum_{\sigma \in S_n} x_{\sigma(n)} R_{n-1}(x_{\sigma(1)}, \dots, x_{\sigma(n-1)}) \exp(2\pi i [\sum_{i < j < n} \langle v_{\sigma(i)}, v_{\sigma(j)} \rangle + \langle \sum_{j \neq m} v_j, v_m \rangle]) &= \\ \sum_{m=1}^n x_m R_{\bar{v}^{(m)}}(x_1, \dots, \widehat{x_m}, \dots, x_n) \exp(2\pi i \langle \sum_{j \neq m} v_j, v_m \rangle). \end{aligned}$$

If $\theta = 0$ or if vectors $v_1, \dots, v_n$ belong to a rank-1 subgroup $\mathbb{Z} \subset \mathbb{Z}^2$ then we have

$$R_{\overline{v}}(x) = \sum_{\sigma \in S_n} R_n(x^\sigma).$$

Applying Proposition 1.7 we deduce that this function has no poles. Since it is homogeneous of degree 0 it must be a constant. To compute this constant we substitute $x_1 = \dots = x_n = 1$ and obtain that in this case $R_{\overline{v}}(x) \equiv 1$. This immediately implies (ii).

To prove part (iii) it suffices to check (1.5) as a formal identity in $A_\theta[[z, t]]$. We claim that both sides solve the same differential equation $\frac{\partial f}{\partial z}(z, t) = tf(z, t)$ with the same initial condition $f(0, t) = \text{Exp}_l(\tau, a)$. Indeed, this is clear for the right-hand side of (1.5), so we just have to check this for the left-hand side. It is easy to check that

$$\frac{1}{D!} \cdot \frac{\partial[(a+z)_D]}{\partial z} = \frac{(a+z)_{D-(0)}}{(D-(0))!}$$

if $D - (0) \geq 0$, otherwise the derivative is zero. This implies that the derivative of $\text{Exp}_l(z+a)$ with respect to z is equal to $t \text{Exp}_l(z+a)$ as we claimed. $\square$

Remark. For n lattice vectors $(v_1, \dots, v_n)$ such that $v_1 + \dots + v_n = 0$ we obtain from the above proof that

$$\sum_{m=1}^n \iota(v_m) f_{n-1}^*(v_1, \dots, \widehat{v_m}, \dots, v_n) = 0.$$

If $v_1, \dots, v_n$ are sufficiently generic this reduces to the identity of Corollary 1.6.

Corollary 1.11. *If $\text{Exp}_l(\tau, a) = \text{Exp}_l(\tau, b)$ for some $a, b \in A_\theta[[t]]$ then $a = b$.*

Proof. Using part (i) of the above theorem we immediately deduce that $b = a + z$ for some $z \in \mathbb{C}[[t]]$. Now part (iii) implies that $\exp(tz) = 1$ and hence $z = 0$. $\square$

We can also define the right exponential map $\text{Exp}_r(\tau, \cdot)$ by setting

$$\text{Exp}_r(\tau, a) = \text{Exp}_l(\overline{\tau}, a^*)^*.$$

Then

$$\delta_\tau(\text{Exp}_r(\tau, a)) = t\delta_\tau(a) \cdot \text{Exp}_r(\tau, a).$$

Using the definition of $\text{Exp}_l(\tau, \cdot)$ we get

$$\text{Exp}_r(\tau, a) = \sum_{D \in \text{Div}(\mathbb{Z}^2)_{\geq 0}} \frac{t^{\deg(D)}}{D!} c_{-\theta}^*(D) a_D U_{s(D)}.$$

Proposition 1.12. *For every $a \in A_\theta[[t]]$ one has*

$$\text{Exp}_l(\tau, a)^{-1} = \text{Exp}_r(\tau, -a).$$

Proof. It suffices to prove that for every divisor $D > 0$ on $\mathbb{Z}^2$ one has

$$\sum_{D_1, D_2 \in \text{Div}(\mathbb{Z}^2)_{\geq 0}: D_1 + D_2 = D} \frac{(-1)^{\deg D_2}}{D_1! D_2!} c_\theta^*(D_1) c_{-\theta}^*(D_2) \exp(2\pi i \langle s(D_1), s(D_2) \rangle_\theta) = 0.$$

This is equivalent to the following identity that should hold for every n -tuple $(v_1, \dots, v_n)$ of lattice vectors (where $n > 0$):

$$\sum_{I=\{i_1, \dots, i_r\} \subset [1, n]} (-1)^{n-r} f_r^*(v_{i_1}, \dots, v_{i_r}; \theta) f_{n-r}^*(v_{j_1}, \dots, v_{j_{n-r}}; -\theta) \\ \times \exp(2\pi i \langle v_{i_1} + \dots + v_{i_r}, v_{j_1} + \dots + v_{j_{n-r}} \rangle_\theta) = 0,$$

where the sum is taken over all subsets $I = \{i_1, \dots, i_r\}$ of $[1, n] = \{1, 2, \dots, n\}$ and we denote by $\{j_1, \dots, j_{n-r}\}$ the complement to I . Recalling the definition of the functions f_n^* we see that it suffices to prove the following identity between rational functions of $x_1, \dots, x_n$:

$$\sum_{r=0}^n \sum_{\sigma \in S_n} (-1)^{n-r} R_r(x_{\sigma(1)}, \dots, x_{\sigma(r)}) R_{n-r}(x_{\sigma(n)}, \dots, x_{\sigma(r+1)}) \\ \exp(2\pi i [\sum_{i < j \leq r} \langle v_{\sigma(i)}, v_{\sigma(j)} \rangle_\theta + \sum_{i > j > r} \langle v_{\sigma(i)}, v_{\sigma(j)} \rangle_{-\theta} + \langle \sum_{i \leq r} v_{\sigma(i)}, \sum_{j > r} v_{\sigma(j)} \rangle_\theta]) = 0.$$

Due to the skew-symmetric of $\langle \cdot, \cdot \rangle_\theta$ we have

$$\langle v, v' \rangle_{-\theta} = \langle v', v \rangle_\theta.$$

Using this observation we can rewrite our identity as

$$\sum_{r=0}^n \sum_{\sigma \in S_n} (-1)^{n-r} R_r(x_{\sigma(1)}, \dots, x_{\sigma(r)}) R_{n-r}(x_{\sigma(n)}, \dots, x_{\sigma(r+1)}) \exp(2\pi i \sum_{i < j} \langle v_{\sigma(i)}, v_{\sigma(j)} \rangle_\theta) = 0.$$

Thus, it is enough to prove that

$$\sum_{r=0}^n (-1)^{n-r} R_r(x_1, \dots, x_r) R_{n-r}(x_n, \dots, x_{r+1}) = 0.$$

This is equivalent to the following identity:

$$\sum_{i=0}^{n-1} \frac{(-1)^i}{x_1(x_1 + x_2) \dots (x_1 + \dots + x_{n-i}) x_n(x_n + x_{n-1}) \dots (x_n + \dots + x_{n-i+1})} = \\ \frac{(-1)^{n-1}}{x_n(x_n + x_{n-1}) \dots (x_n + \dots + x_1)}.$$

We claim that more generally for $0 < j < n$ one has

$$\sum_{i=0}^j \frac{(-1)^i}{x_1(x_1 + x_2) \dots (x_1 + \dots + x_{n-i}) x_n(x_n + x_{n-1}) \dots (x_n + \dots + x_{n-i+1})} = \\ \frac{(-1)^j}{x_1(x_1 + x_2) \dots (x_1 + \dots + x_{n-j-1}) \cdot (x_1 + \dots + x_n) \cdot x_n(x_n + x_{n-1}) \dots (x_n + \dots + x_{n-j+1})}.$$

This can be easily checked by induction in j . $\square$

Corollary 1.13. *One has*

$$\text{tr}(\text{Exp}_l(\tau, a)) \cdot \text{tr}(\text{Exp}_r(\tau, -a)) = \mathbf{s}(\tau, a)^{-1}.$$

Proof. This follows from the relation $\text{Exp}_l(\tau, a) = \text{tr}(\text{Exp}_l(\tau, a))\mathbf{E}_l(\tau, a)$ and the similar relation for right exponentials. $\square$

Finally, let us state the analogue of the equation $\exp(x)\exp(y) = \exp(x+y)$ for our exponentials.

Proposition 1.14. *For every $a, b \in A_\theta[[t]]$ there exists a unique $\varphi(a, b) \in A_\theta[[t]]$ such that*

$$\text{Exp}_l(a) \text{Exp}_l(b) = \text{Exp}_l(\varphi(a, b) + b) \quad (1.6)$$

and

$$\delta_\tau(\varphi(a, b)) = \text{Exp}_l(\tau, b)^{-1} \delta_\tau(a) \text{Exp}_l(\tau, b).$$

Proof. Since the element $\text{Exp}_l(\tau, b)^{-1} \delta_\tau(a) \text{Exp}_l(\tau, b) \in A_\theta[[t]]$ has zero trace we can find some element $\varphi' \in A_\theta[[t]]$ such that

$$\delta_\tau(\varphi') = \text{Exp}_l(\tau, b)^{-1} \delta_\tau(a) \text{Exp}_l(\tau, b).$$

Now one can easily check that the left-hand side of (1.6) is a solution of the equation

$$\delta_\tau(x) = tx\delta_\tau(\varphi' + b).$$

It follows that

$$\text{Exp}_l(a) \text{Exp}_l(b) = f(t) \text{Exp}_l(\varphi' + b)$$

for some $f(t) \in 1 + t\mathbb{C}[[t]]$. Writing $f(t)$ in the form $\exp(tz)$ for $z \in \mathbb{C}[[t]]$ and using part (iii) of Theorem 1.10 we conclude that (1.6) holds for $\varphi(a, b) = z + \varphi'$. The uniqueness follows from Corollary 1.11. $\square$

1.5. Compatibility with $\text{SL}_2(\mathbb{Z})$ -action. The group $\text{SL}_2(\mathbb{Z})$ acts on the algebra A_θ by automorphisms. Namely, to an element $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$ one associates an automorphism

$$\alpha_g : A_\theta \rightarrow A_\theta : U_v \mapsto U_{gv}.$$

These automorphisms are compatible with derivations δ_τ and the modular action of $\text{SL}_2(\mathbb{Z})$ on τ :

$$\alpha_g \delta_\tau \alpha_g^{-1} = \delta_{g\tau}$$

where $g\tau = (a\tau + b)/(c\tau + d)$.

The only way in which the standard basis of $\mathbb{Z}^2$ enters into our formulae for $\mathbf{E}_l(\tau, a)$ and $\text{Exp}_l(\tau, a)$ is through the definition of the homomorphism $\iota = \iota_\tau : \mathbb{Z}^2 \rightarrow \mathbb{C}$. For an element $g \in \text{SL}_2(\mathbb{Z})$ we have

$$\iota_{g\tau}(v) = \iota_\tau({}^t g v) \cdot j_\tau(g)^{-1},$$

where ${}^t g$ is the transpose of g , and $j_\tau(g) = c\tau + d$. Since the functions $f_n^*(v_1, \dots, v_n)$ (resp., $f_n(v_1, \dots, v_n)$) are homogeneous of degree 0 this implies that

$$f_n^*(v_1, \dots, v_n; g\tau) = f_n^*({}^t g v_1, \dots, {}^t g v_n; \tau)$$

and similarly for the functions f_n . This easily implies the following result.

Proposition 1.15. *For every $a \in A_\theta[[t]]$ one has*

$$\begin{aligned}\mathbf{E}_l(g'\tau, \alpha_g(a)) &= \alpha_g \mathbf{E}_l(\tau, a), \\ \mathbf{E}_r(g'\tau, \alpha_g(a)) &= \alpha_g \mathbf{E}_r(\tau, a), \\ \text{Exp}_l(g'\tau, \alpha_g(a)) &= \alpha_g \text{Exp}_l(\tau, a), \\ \text{Exp}_r(g'\tau, \alpha_g(a)) &= \alpha_g \text{Exp}_r(\tau, a), \\ \mathbf{s}(g'\tau, \alpha_g(a)) &= \mathbf{s}(\tau, a),\end{aligned}$$

where $g' = {}^t g^{-1}$.

2. CONVERGENT EXPONENTIALS, LOGARITHMIC DERIVATIVES AND THE DISCRIMINANT HYPERSURFACE

2.1. Norms. In this section we establish notation for some of the norms we use on A_θ .

First of all we have the operator algebra norm $\|\cdot\|$ on A_θ : $\|a\|$ is defined to be the norm of the operator of multiplication by a on the L^2 -completion of A_θ (it does not matter whether one considers left or right multiplication). Recall that the C^* -algebra $\overline{A}_\theta$ is defined as the completion of A_θ with respect to this norm.

We will also consider the L^2 -norm

$$\|a\|_0 = \left(\sum_{v \in \mathbb{Z}^2} |a_v|^2 \right)^{1/2},$$

where $a = \sum_{v \in \mathbb{Z}^2} a_v U_v$. By the definition, for every $a, b \in A_\theta$ we have

$$\|ab\|_0 \leq \|a\| \cdot \|b\|_0.$$

For every $s \geq 0$ we can also consider the norm

$$\|a\|_s^2 = \sum_{i=0}^s \|\delta_\tau^i a\|_0^2.$$

The completion of A_θ with respect to this norm is the *Sobolev space* W_s . We have a sequence of embeddings $W_0 \supset W_1 \supset W_2 \supset \dots$. These spaces enjoy all the usual properties of the Sobolev spaces (see [4]). In particular, it can be extended to a sequence of spaces (W_s) , where $s \in \mathbb{Z}$, so that W_s is dual to W_{-s} . Also, the intersection of all W_s is A_θ and the topology on A_θ is the one determined by the collection of seminorms coming from the W_s .

Finally, at one point we will use the norm

$$\|a\|_{l^1} := \sum_{v \in \mathbb{Z}^2} |a_v|.$$

For $a, b \in A_\theta$ we have

$$\|ab\|_{l^1} \leq \sum_{v \in \mathbb{Z}^2} \|a_v U_v b\|_{l^1} = \sum_{v \in \mathbb{Z}^2} |a_v| \cdot \|b\|_{l^1} = \|a\|_{l^1} \cdot \|b\|_{l^1}.$$

On the other hand,

$$\|a\| \leq \sum_{v \in \mathbb{Z}^2} \|a_v U_v\| = \|a\|_{l^1}.$$

2.2. Convergence criteria. If one wants to solve the equation

$$\delta_\tau(x) = x\delta_\tau(a) \quad (2.1)$$

for given $a \in A_\theta$ one can try to consider the series

$$x = \mathbf{E}_l(\tau, a)|_{t=1} = 1 + a_1 + a_2 + \dots \quad (2.2)$$

obtained by substituting $t = 1$ into $\mathbf{E}_l(\tau, a) = 1 + ta_1 + t^2a_2 + \dots$. This series does not always converge even for $\theta = 0$ (remember that in the commutative case it represents the function $\exp(ta)/\text{tr}(\exp(ta))$). One of our goals in this section is to show that (2.2) always converges for sufficiently small a .

Remark. In the commutative case the convergence improves when we switch from $\mathbf{E}_l(\tau, a)$ to $\text{Exp}_l(\tau, a)$ since the latter series becomes the usual exponential. However, this does not seem to work in noncommutative case. Indeed, as we will see in Proposition 2.13(ii) there exists $a \in A_\theta$ for which (2.1) has no invertible solutions. By Proposition 1.12 this means that either $\text{Exp}_l(\tau, a)|_{t=1}$ or $\text{Exp}_r(\tau, -a)|_{t=2}$ diverges.

The next lemma follows from Corollary 7.16 of [10]. Since our particular case is much easier we present here a more direct proof due to Chris Phillips.

Lemma 2.1. *If $x \in A_\theta$ is invertible as an element of $\overline{A}_\theta$ then $x \in A_\theta^*$.*

Proof. First, we claim that for every $a \in A_\theta$ such that $\|a\| < 1$, the series $1 + a + a^2 + \dots$ converges to an element of A_θ . Indeed, it suffices to show that it converges absolutely in every Sobolev space W_s . But this follows from the fact that for every k there exists a constant C (depending on a) such that $\|\delta^k(a^n)\| \leq C\|a\|^{n-k}$ for $n > k$. Thus, the assertion is true when $\|x - 1\| < 1$. The general case can be reduced to this by the following trick. Let x be an element of A_θ such that there exists an inverse $x^{-1} \in \overline{A}_\theta$. Choose $y \in A_\theta$ such $\|y - x^{-1}\| < \|x\|^{-1}$. Then the element $xy \in A_\theta$ satisfies $\|xy - 1\| < 1$. Hence, $xy \in A_\theta^*$. It follows that x has a right inverse in A_θ . Similar argument shows that $yx \in A_\theta^*$ and hence x has a left inverse. This implies that $x \in A_\theta^*$. $\square$

Remark. In fact, it is also true that if $xy = 1$ in A_θ then $yx = 1$. This follows from the existence of a finite positive faithful trace on A_θ (cf. [1], second half of p. 101 and Exer. 6 and 9 of ch. 4). Here is another argument using Rieffel's classification of finitely generated projective A_θ -modules in [8] (we assume that θ is irrational). Let $I \subset A_\theta$ the set of all a such that $xa = 0$. Then I is a right ideal and we have $A_\theta = I \oplus yA_\theta$. It follows that yA_θ and I are projective modules. Furthermore, yA_θ is isomorphic to A_θ hence $\text{rk } yA_\theta = 1$. It follows that $\text{rk } I = 0$, so $I = 0$.

Theorem 2.2. *Let*

$$d = \min_{(m,n) \in \mathbb{Z}^2 \setminus \{(0,0)\}} |m\tau + n|.$$

Then for every $a \in A_\theta$ with $\text{tr}(a) = 0$ such that $\|\delta_\tau(a)\| < 2\pi d$, the series (2.2) converges to an element $\mathbf{e}_l(\tau, a) \in A_\theta$ satisfying the equation

$$\delta_\tau(\mathbf{e}_l(\tau, a)) = \mathbf{e}_l(\tau, a)\delta_\tau(a).$$

Furthermore, if $\|\delta_\tau(a)\|_{l^1} < \pi d$ then $\mathbf{e}_l(\tau, a) \in A_\theta^*$. The map $a \mapsto \mathbf{e}_l(\tau, a)$ from $U_\tau = \{a \in H : \|\delta_\tau(a)\|_{l^1} < \pi d\}$ to A_θ^* is continuous.

Proof. Recall that a_i 's are defined inductively by the conditions $\delta(a_i) = a_{i-1}\delta(a)$, $\text{tr}(a_i) = 0$, where $\delta = \delta_\tau$. It follows that

$$\|a_i\|_0 \leq (2\pi d)^{-1} \|\delta(a_i)\|_0 \leq (2\pi d)^{-1} \|\delta(a)\| \cdot \|a_{i-1}\|_0.$$

Hence, $\|a_n\|_0 \leq (\frac{\|\delta(a)\|}{2\pi d})^n$ for every $n \geq 0$. It follows that the series $1 + a_1 + a_2 + \dots$ converges to an element x in the L^2 -completion of A_θ . Furthermore, since

$$\|\delta(a_n)\|_0 \leq \|\delta(a)\| \cdot \|a_{n-1}\|_0 \leq \|\delta(a)\| \cdot (\frac{\|\delta(a)\|}{2\pi d})^{n-1},$$

we see that in fact x belongs to the Sobolev space W_1 and satisfies the equation

$$\delta(x) = x\delta(a).$$

This implies that $x \in A_\theta$.

To prove the last assertion we first observe that $\|a_n\|_{l^1} \leq r^n$ for all n , where $r = \|\delta(a)\|_{l^1}/(2\pi d)$ (this is deduced in the same way as above). Hence,

$$\|a_1 + a_2 + \dots\| \leq \|a_1 + a_2 + \dots\|_{l^1} \leq \frac{r}{1-r} < 1$$

if $r < 1/2$. Therefore, x is invertible as an element of $\overline{A}_\theta$. It remains to apply Lemma 2.1. $\square$

The following result shows that our exponentials converge if all the coefficients of a belong to closed halfplanes not containing zero.

Proposition 2.3. *Let $a = \sum_v a_v U_v \in A_\theta$ be an element with $\text{tr}(a) = 0$. Assume that there exists a homomorphism $h : \mathbb{Z}^2 \rightarrow \mathbb{R}$ and a positive constant ϵ such that $h > \epsilon$ on $\text{supp}(a) = \{v \in \mathbb{Z}^2 : a_v \neq 0\}$. Then the above series $1 + a_1 + a_2 + \dots$ converges to an element of A_θ^* .*

Proof. Since the elements $a_i \in H$ are defined inductively by $a_1 = a$ and $\delta(a_i) = a_{i-1}\delta(a)$, we deduce that $h > n\epsilon$ on $\text{supp}(a_n)$. Let $d_n = \min_{(m,n): h(m,n) > n\epsilon} |m\tau + n|$. Then we have

$$\|a_n\|_0 \leq (2\pi d_n)^{-1} \|\delta(a_n)\|_0 \leq (2\pi d_n)^{-1} \|\delta(a)\| \cdot \|a_{n-1}\|_0.$$

Since d_n grows linearly with n this immediately implies convergence of the series $\sum a_n$ in the L^2 -norm. Arguing as in the proof of Theorem 2.2 we derive that $x = 1 + a_1 + a_2 + \dots$ belongs to A_θ^* . $\square$

Remark. In the situation of the above proposition one has $\text{Exp}_l(\tau, a)|_{t=1} = \mathbf{E}_l(\tau, a)|_{t=1}$. However, at present we do not know any criteria for convergence of $\text{Exp}_l(\tau, a)|_{t=1}$ similar to Theorem 2.2.

2.3. Discriminant hypersurface. Recall that $H \subset A_\theta$ is the set of elements a such that $\text{tr}(a) = 0$. For every $a \in A_\theta$ consider the operator

$$d_a := \delta_\tau + \text{ad}(a) : A_\theta \rightarrow A_\theta.$$

We are interested in the following subset of H :

$$\Theta = \Theta_\tau = \{a \in H : \dim \ker(d_a) > 1\}.$$

Note that we always have $\mathbb{C} \subset \ker(d_a)$, so Θ coincides with the set of $a \in H$ such that $\ker(d_a|_H) \neq 0$. We are going to show that Θ_τ is a complex hypersurface in H in the sense that it can be given locally as the zero set of a holomorphic function on H .

Lemma 2.4. *For every $a \in A_\theta$ one has $\ker(d_a) \simeq \text{Hom}_{\mathcal{C}}(E, E)$, $\text{coker}(d_a) \simeq \text{Ext}_{\mathcal{C}}^1(E, E)$, where E is the holomorphic bundle $(A_\theta, \overline{\nabla}_a := \delta_\tau + a)$. In particular, the operator d_a is Fredholm of index zero. The same assertions are true if we replace d_a by its extension to Sobolev spaces $d_a : W_{s+1} \rightarrow W_s$.*

Proof. The identification of $\ker(d_0)$ with endomorphisms of E in the category $\mathcal{C}$ follows from the definitions. The identification of $\text{coker}(d_0)$ with $\text{Ext}_{\mathcal{C}}^1(E, E)$ is constructed similarly to Proposition 2.4 of [6]). Note that by Serre duality $\text{Ext}_{\mathcal{C}}^1(E, E) \simeq \text{Hom}_{\mathcal{C}}(E, E)^*$ (see [4]). The remaining assertions are proved using the same techniques as in Theorem 2.8 of [4]. $\square$

Let $\mathcal{F}_0(H)$ be the space of Fredholm operators of index zero on $W_0(H)$, where $W_0(H) = \{a \in W_0(A_\theta) : \text{tr}(a) = 0\}$. Recall (see [7],[2]) that there is a natural holomorphic line bundle $\mathcal{L}$ over $\mathcal{F}_0(H)$ (called the *determinant line bundle*). By definition, $\mathcal{L}$ is trivial over each open set $\mathcal{U}_A \subset \mathcal{F}_0(H)$, where $A : W_0(H) \rightarrow W_0(H)$ is a fixed trace class operator, $\mathcal{U}_A$ consists of operators T such that $T + A$ is invertible. The transition functions of $\mathcal{L}$ with respect to this open covering are given by the functions

$$g_{A,B}(T) = \det_F(1 + (A - B)(T + B)^{-1}) = \det_F((T + A)(T + B)^{-1}),$$

where A, B are trace class operators, and $\det_F$ denotes the Fredholm determinant.

Let $\mathcal{Z} \subset \mathcal{F}_0(H)$ be the complement of $\mathcal{U}_0$, i.e., the subset consisting of T such that $\ker(T) \neq 0$. Then $\mathcal{Z}$ is a complex hypersurface in $\mathcal{F}_0(H)$. More precisely, there exists a global holomorphic section s of the determinant line bundle $\mathcal{L}$ such that $\mathcal{Z}$ coincides with the zero locus of s . Under the standard trivialization of $\mathcal{L}$ over $\mathcal{U}_A$ the section s corresponds to the holomorphic function

$$s_A(T) = \det_F(1 - (T + A)^{-1}A).$$

One can easily check that $s_B = g_{A,B}s_A$, so these functions glue into a global section of $\mathcal{L}$. One has $s_A(T) = 0$ iff $1 - (T + A)^{-1}A$ has a nonzero kernel. But $1 - (T + A)^{-1}A = (T + A)^{-1}T$, so $s_A(T) = 0$ iff $\ker(T) \neq 0$. This proves that the zero locus of s coincides with $\mathcal{Z}$.

We have a map

$$H \rightarrow \mathcal{F}_0(H) : a \mapsto d_a|_{W_0(H)}.$$

Note that the restriction of d_a to H (resp., $W_0(H)$) still has index zero, since $d_a(A_\theta) \subset H$ and $\ker(d_a) = \mathbb{C} \oplus \ker(d_a|_H)$. Abusing the notation we denote by $\mathcal{L}$ also the pull-back of the determinant line bundle to H under the above map.

Proposition 2.5. (i) The subset $\Theta \subset H$ coincides with the zero locus of the global holomorphic section $s(d_a|_{W_0(H)})$ of $\mathcal{L}$ over H .

(ii) For every one-dimensional subspace $L \subset H$ the intersection $\Theta \cap L$ is a discrete subset of $L \setminus \{0\}$.

Proof. (i) This is clear since Θ is the preimage of $\mathcal{Z} \subset \mathcal{F}_0(H)$.

(ii) We observe that the operator $\delta_\tau : W_0(H) \rightarrow W_{-1}(H)$ is an isomorphism and the operator $\text{ad}(a) : W_0(H) \rightarrow W_{-1}(H)$ is compact for every $a \in A_\theta$ (see the proof of Theorem 2.8 in [4]). Hence, for every $a \in H$ and $t \in \mathbb{C}$ we have $ta \in \Theta$ iff $-t^{-1}$ belongs to the spectrum of the compact operator $\delta_\tau^{-1} \text{ad}(a) : W_0(H) \rightarrow W_0(H)$. $\square$

Remark. Part (ii) of the above proposition implies that Θ contains no linear subspaces of H . On the other hand, we will see later that Θ is swept by infinite-dimensional *affine* subspaces (see Theorem 2.17).

We can also consider more general loci

$$\Theta_\tau^{(n)} = \{a \in H : \dim \ker(d_a) > n + 1\} = \{a \in H : \dim \ker(d_a|_H) > n\}$$

for $n \geq 0$. For $n = 0$ we get $\Theta_\tau^{(0)} = \Theta_\tau$. Since the kernel of d_a coincides with the kernel of its extension to W_0 , it follows that the loci $\Theta_\tau^{(n)}$ are closed.

2.4. Logarithmic derivative and connected components of groups of invertible elements. In this section we will study the relation between holomorphic structures and the group of invertible elements in A_θ .

Theorem 2.6. Consider the map

$$L_\tau : A_\theta^* \rightarrow A_\theta : x \mapsto x^{-1} \delta_\tau(x).$$

(i) The composition

$$\chi_\tau := \text{tr} \circ L_\tau : A_\theta^* \rightarrow \mathbb{C}$$

is a locally constant homomorphism with image $(2\pi i)(\mathbb{Z} + \mathbb{Z}\tau) \subset \mathbb{C}$ and kernel $(A_\theta^*)_0$, the connected component of 1 in A_θ^* .

(ii) Let $H \subset A_\theta$ be the set of elements a such that $\text{tr}(a) = 0$. Let

$$\Omega_\tau = \{a \in H : \dim \ker(\delta_\tau + \text{ad}(a)) = 1\}$$

be the complement to the hypersurface $\Theta_\tau \subset H$. Then Ω_τ is a dense open subset of H and L_τ induces homeomorphisms

$$\begin{aligned} A_\theta^*/\mathbb{C}^* &\xrightarrow{\sim} \Omega_\tau + 2\pi i(\mathbb{Z} + \mathbb{Z}\tau), \\ (A_\theta^*)_0/\mathbb{C}^* &\xrightarrow{\sim} \Omega_\tau. \end{aligned}$$

Proof. From the Leibnitz rule we immediately get

$$L_\tau(xy) = y^{-1} L_\tau(x)y + L_\tau(y). \quad (2.3)$$

Taking traces we see that $\chi_\tau = \text{tr} \circ L_\tau$ is a homomorphism. It is easy to see that the derivative of L_τ at the point $x \in A_\theta^*$ is $a \mapsto x^{-1} \delta_\tau(ax^{-1})x$. Hence, the derivative of χ_τ is $a \mapsto \text{tr}(x^{-1} \delta_\tau(ax^{-1})x) = 0$. This implies that χ_τ is locally constant. In particular, $\chi_\tau((A_\theta^*)_0) = 0$, i.e., $L_\tau((A_\theta)_0) \in H$.

Next we claim that the map $L_\tau : (A_\theta^*)_0/\mathbb{C}^* \rightarrow H$ is a local homeomorphism. Indeed, (2.3) shows that it is enough to check that L_τ is a local homeomorphism in a neighborhood of $1 \in (A_\theta^*)_0/\mathbb{C}^*$. Then we can use the map $\mathbf{e}_l$ constructed in Theorem 2.2 to get a local inverse to L_τ .

We know that the space of holomorphic endomorphisms of $(A_\theta, \delta_\tau + a)$ can be identified with $\ker(\delta_\tau + \text{ad}(a))$ (see Lemma 2.4). Using the classification of holomorphic bundles on noncommutative tori given in [4] we derive that the dimension of this space is 1 iff there exists a holomorphic isomorphism of $(A_\theta, \overline{\nabla})$ with a standard holomorphic bundle $(A_\theta, \delta_\tau + z)$ for some $z \in \mathbb{C}$. In other words, $\ker(\delta_\tau + \text{ad}(a)) = \mathbb{C}$ iff there exists $x \in A_\theta^*$ such that $a \equiv L_\tau(x) \bmod \mathbb{C}$. Thus, the map

$$A_\theta^*/\mathbb{C}^* \rightarrow H : x \mapsto L_\tau(x) - \text{tr}(L_\tau(x)) \quad (2.4)$$

has Ω_τ as an image. As we have seen above this map is a local homeomorphism, so Ω_τ is open in H . Now we claim that nonempty fibers of (2.4) are exactly orbits of the action of the central subgroup $\mathbb{Z}^2 \subset A_\theta^*/\mathbb{C}^*$. Indeed, assume that $L_\tau(x) = L_\tau(y) \bmod \mathbb{C}$ for some $x, y \in A_\theta^*$. Set $z = xy^{-1}$. Then

$$L_\tau(x) = L_\tau(z)y = y^{-1}L_\tau(z)y + L_\tau(y)$$

which implies that $y^{-1}L_\tau(z)y \in \mathbb{C}$, hence, $L_\tau(z) \in \mathbb{C}$. But this is possible only if z is proportional to U_v for some $v \in \mathbb{Z}^2$. This proves our claim about the fibers of the map (2.4).

Proposition 2.5(ii) immediately implies that Ω_τ is connected (and dense in H). Since we have identified the quotient of $A_\theta^*/\mathbb{C}^*$ by $\mathbb{Z}^2$ with Ω_τ , it follows that the embedding $\mathbb{Z}^2 \subset A_\theta^*$ induces an isomorphism on connected components. Since χ_τ is locally constant this implies that

$$\chi_\tau(A_\theta^*) = \chi_\tau(\mathbb{Z}^2) = (2\pi i)(\mathbb{Z} + \mathbb{Z}\tau).$$

We also see that $L_\tau : (A_\theta^*)_0/\mathbb{C}^* \rightarrow \Omega_\tau$ is a homeomorphism. $\square$

Corollary 2.7. *For every $a \in H$ there exists a nonzero $x \in A_\theta$ such that $\delta_\tau(x) = ax$ (resp., $\delta_\tau(x) = xa$).*

Proof. If $-a \in \Omega_\tau$ then by the above theorem there exists $x \in A_\theta^*$ such that $\delta_\tau(x) = -xa$. Therefore, $\delta_\tau(x^{-1}) = ax^{-1}$. If $-a \notin \Omega_\tau$ then the holomorphic bundle $(A_\theta, \overline{\nabla}_{-a} = \delta_\tau - a)$ is decomposable. Since one of the indecomposable factors should be of positive degree this implies that $H^0(A_\theta, \overline{\nabla}_{-a}) \neq 0$. The case of the equation $\delta_\tau(x) = xa$ reduces to the previous case by using the identification $A_\theta^{opp} = A_{-\theta}$. $\square$

As another corollary we get a new proof of the following well known fact.

Corollary 2.8. *The embedding $\mathbb{Z}^2 \subset A_\theta^*$ induces an isomorphism on connected components.*

Using the results of [4] we can give one more characterization of the open subset $\Omega_\tau \subset H$. As in section 2.3 this leads to a set of local equations for its complement Θ_τ . For every $a \in H$ let us set $\overline{\nabla}_a = \delta_\tau + a$.

Proposition 2.9. *For $a \in H$ one has $a \in \Omega_\tau$ iff there exists $z \in \mathbb{C}$ such that*

$$\ker(\overline{\nabla}_{a+z}) = \{x : \delta_\tau(x) + (a+z)x = 0\}$$

is zero. Moreover, for $a \in \Omega_\tau$ this holds for all $z \notin 2\pi i(\mathbb{Z} + \mathbb{Z}\tau)$. Hence, if we choose any $z_0 \notin 2\pi i(\mathbb{Z} + \mathbb{Z}\tau)$ then $a \in \Omega_\tau$ iff the operator $\overline{\nabla}_{a+z_0}$ is invertible.

Proof. If $a \in \Omega_\tau$ then this follows from the fact that cohomology of the standard holomorphic bundle $(A_\theta, \delta_\tau + z)$ vanishes for $z \notin 2\pi i(\mathbb{Z} + \mathbb{Z}\tau)$. If $a \notin \Omega_\tau$ then it suffices to observe that the holomorphic bundle $(A_\theta, \overline{\nabla}_{a+z})$ has a direct factor of positive degree. $\square$

One can generalize some of the assertions of Theorem 2.6 to a slightly more general context.

Proposition 2.10. *Let $(E, \overline{\nabla})$ be a basic right module over A_θ equipped with a holomorphic structure. Then the map*

$$\chi_E : \text{Aut}_{A_\theta}(E) \rightarrow \mathbb{C} : x \mapsto \text{tr}(x^{-1}[\overline{\nabla}, x])$$

is a locally constant homomorphism that does not depend on $\overline{\nabla}$. Its kernel coincides with the connected component of 1 in $\text{Aut}_{A_\theta}(E)$ and its image is the lattice $\frac{2\pi i}{\text{rk}(E)}(\mathbb{Z} + \tau\mathbb{Z}) \subset \mathbb{C}$, where $\text{rk}(E) \in (\mathbb{Z} + \mathbb{Z}\theta) \cap \mathbb{R}_{>0}$ is the rank of E . If $E = E' \oplus E''$ is a decomposition into the direct sum of A_θ -modules, where E' is also basic, then one has the commutative diagram

$$\begin{array}{ccc} \text{Aut}_{A_\theta}(E') & \xrightarrow{\chi_{E'}} & \mathbb{C} \\ \downarrow & & \downarrow \text{rk}(E')/\text{rk}(E) \\ \text{Aut}_{A_\theta}(E) & \xrightarrow{\chi_E} & \mathbb{C} \end{array} \quad (2.5)$$

Proof. Any other holomorphic structure on E has form $\overline{\nabla} + \phi$ for some $\phi \in \text{End}_{A_\theta}(E)$. But

$$\text{tr}(x^{-1}[\overline{\nabla} + \phi, x]) = \text{tr}(x^{-1}[\overline{\nabla}, x]) + \text{tr}(x^{-1}\phi x - \phi) = \text{tr}(x^{-1}[\overline{\nabla}, x]).$$

This shows that χ_E is independent of $\overline{\nabla}$. If we choose $\overline{\nabla}$ to be standard (see [4]) then the pair $(\text{End}_{A_\theta}, \text{ad } \overline{\nabla})$ can be identified with a pair of the form $(A_{\theta'}, \delta_\tau/r)$ for some θ' , where $r = \text{rk}(E)$ (see [6], Prop. 2.1). It remains to apply Theorem 2.6.

Since the homomorphism χ_E does not depend on a choice of a holomorphic structure we can choose a holomorphic structure on E compatible with the decomposition $E = E' \oplus E''$. Then commutativity of (2.5) follows from the compatibility of the embedding $i : \text{End}_{A_\theta}(E') \rightarrow \text{End}_{A_\theta}(E)$ with normalized traces:

$$\text{tr}(i(x)) = \frac{\text{rk}(E')}{\text{rk}(E)} \text{tr}(x).$$

$\square$

As a corollary we get a new proof of the following well known fact.

Corollary 2.11. *Let E be a basic right module over A_θ , and let $E = E' \oplus E''$ be a decomposition into the direct sum of A_θ -modules, where E' is also a basic module. Then the natural homomorphism $\text{Aut}_{A_\theta}(E') \rightarrow \text{Aut}_{A_\theta}(E)$ induces an isomorphism on connected components.*

Proof. Apply commutative diagram (2.5) together with the fact that the image of χ_E (resp., of $\chi_{E'}$) is $2\pi i \text{rk}(E)(\mathbb{Z} + \mathbb{Z}\tau)$ (resp., $2\pi i \text{rk}(E')(\mathbb{Z} + \mathbb{Z}\tau)$). $\square$

Corollary 2.12. *Let E be a basic right module over A_θ , and let $E = E' \oplus E''$ be a decomposition into a direct sum of A_θ -modules. Then the natural homomorphism $\text{Aut}_{A_\theta}(E') \rightarrow \text{Aut}_{A_\theta}(E)$ induces a surjection on connected components.*

Proof. It suffices to choose any decomposition $E' = E'_1 \oplus E'_2$ with E'_1 basic and apply Corollary 2.11. $\square$

Remark. The well known fact that the embedding of A_θ into $\text{Mat}_n(A_\theta)$ induces an isomorphism on connected components of groups of invertible elements (for $\overline{A}_\theta$ this statement is Theorem 8.3 of [8], the case of A_θ follows using its invariance under the holomorphic functional calculus) allows to deduce from Corollary 2.11 the same statement for E' not necessarily basic.

2.5. More on discriminant hypersurface. It would be interesting to study intersections of Θ_τ with finite-dimensional subspaces of H . In the following proposition we consider simplest examples.

Proposition 2.13. (i) *Let $h : \mathbb{Z}^2 \rightarrow \mathbb{R}$ be a homomorphism. Assume that $a \in H$ is an element such that $h > \epsilon$ on $\text{supp}(a)$ for some $\epsilon > 0$. Then $\mathbb{C}a \in \Omega_\tau$. The same conclusion holds if we assume that h takes values in $\mathbb{Q}$ and that $h \geq 0$ on $\text{supp}(a)$.*
(ii) *Let $e \in A_\theta$ be a nontrivial idempotent such that δ_τ preserves $eA_\theta \subset A_\theta$, i.e., $e\delta_\tau(e) = \delta_\tau(e)$. Then $\Theta_\tau \cap \mathbb{C}\delta_\tau(e) = \{\delta_\tau(e)\}$.*

We need the following simple lemma for the proof.

Lemma 2.14. *Let $a = x_0^{-1}\delta_\tau(x_0)$ for some $x_0 \in A_\theta^*$. Then every $x \in A_\theta$ such that $\delta_\tau(x) = xa$ is proportional to x_0 .*

Proof. Indeed, we have

$$\delta(xx_0^{-1}) = \delta(x)x_0^{-1} - xx_0^{-1}\delta(x_0)^{-1}x_0^{-1} = 0,$$

hence xx_0^{-1} is a constant. $\square$

Proof of Proposition 2.13. (i) It suffices to show that the equation $L_\tau(x) = a$ has a solution with $x \in A_\theta^*$. If $h > \epsilon$ on $\text{supp}(a)$ then this follows from Proposition 2.3. If h takes values in $\mathbb{Q}$ and $h \geq 0$ on $\text{supp}(a)$ then we can write $a = b + c$, where $h = 0$ on $\text{supp}(b)$ and $h > \epsilon > 0$ on $\text{supp}(c)$. Since the subalgebra of elements supported on $h = 0$ is commutative and stable under δ_τ , we have $b = L_\tau(x)$ for some $x \in A_\theta^*$ such that $h = 0$

on $\text{supp}(x)$. Now let us consider the element $c' = xcx^{-1}$. Note that $h > 0$ on $\text{supp}(c')$. Therefore, by Proposition 2.3 there exists an element $y \in A_\theta^*$ such that $c' = L_\tau(y)$. Then

$$L_\tau(yx) = x^{-1}L_\tau(y)x + L_\tau(x) = x^{-1}c'x + b = c + b = a.$$

(ii) For $\lambda \in \mathbb{C}^*$ consider the element $x_\lambda = \lambda e + (1 - e) \in A_\theta$. Then $x_\lambda \in (A_\theta^*)_0$ and $x_\lambda^{-1} = \lambda^{-1}e + (1 - e)$. Hence,

$$L_\tau(x_\lambda) = (\lambda^{-1}e + (1 - e))(\lambda - 1)\delta_\tau(e) = (1 - \lambda^{-1})\delta_\tau(e)$$

It follows that $z\delta_\tau(e) \in \Omega_\tau$ for all $z \in \mathbb{C} \setminus \{1\}$. Now let us prove that $\delta_\tau(e) \notin \Omega_\tau$. Assume that there exists $x_0 \in A_\theta^*$ such that

$$x_0^{-1}\delta_\tau(x_0) = \delta_\tau(e).$$

Since we also have $\delta_\tau(e) = e\delta_\tau(e)$, applying Lemma 2.14 to $x = e$ we derive that e should be proportional to x_0 . Since e is not invertible we get a contradiction. $\square$

Remark. There are plenty of idempotents $e \in A_\theta$ such that eA_θ is preserved by δ_τ . In fact, for every number $0 < m\theta + n < 1$ such that $m, n \in \mathbb{Z}$ and $m < 0$ there exists an idempotent e as above with $\text{tr}(e) = m\theta + n$ (see [5]).

Next, we are going to describe the decomposition discriminant locus $\Theta_\tau \subset H$ corresponding to types of idempotents in A_θ .

Lemma 2.15. *Assume that θ is irrational. Then two idempotents e and e' in A_θ are conjugate by an element in $(A_\theta^*)_0$ iff $\text{tr}(e) = \text{tr}(e')$.*

Proof. The “only if” part is trivial. Assume that $\text{tr}(e) = \text{tr}(e')$. Then by Rieffel’s classification of projective A_θ -modules (see [8]) there exist isomorphisms of right A -modules $eA \simeq e'A$ and $(1 - e)A \simeq (1 - e')A$. Therefore, there exists an element $x \in A_\theta^*$ such that $xeA = e'A$ and $x(1 - e)A = (1 - e')A$. This immediately implies the equality of the idempotents $xe x^{-1} = e'$. It remains to show that x can be chosen in $(A_\theta^*)_0$. But this follows easily from Corollary 2.12. $\square$

For every idempotent e let us denote by $A_{e,\tau}$ the set of $a \in A_\theta$ such that the operator $x \mapsto \delta(x) + ax$ preserves the decomposition $A_\theta = eA_\theta \oplus (1 - e)A_\theta$, i.e.,

$$A_{e,\tau} = \{a \in A_\theta \mid \delta(e) + ae \in eA_\theta, \delta(1 - e) + a(1 - e) \in (1 - e)A_\theta\}.$$

Equivalently, $a \in A_{e,\tau}$ iff a satisfies the equations

$$e\delta(e) = ea(1 - e), \quad (1 - e)ae = (1 - e)\delta(1 - e).$$

This shows that $A_{e,\tau}$ is an affine subspace in A_θ with the associated linear subspace $eAe + (1 - e)A(1 - e)$.

Lemma 2.16. *For $x \in A_\theta^*$ and for an idempotent $e \in A_\theta$ the map $a \mapsto x^{-1}ax + L_\tau(x)$ induces a bijection from $A_{e,\tau}$ to $A_{x^{-1}ex,\tau}$.*

Proof. In fact, the natural action of A_θ^* on operators of the form $\delta_\tau + a$ is equivalent to the following action of A_θ^* on A_θ :

$$x * a = xax^{-1} + L_\tau(x^{-1}),$$

where $x \in A_\theta^*$, $a \in A_\theta$. It is clear from the definition that the action of x^{-1} sends $A_{e,\tau}$ to $A_{x^{-1}ex,\tau}$. $\square$

We refer to the action of A_θ^* on A_θ introduced above as *twisted action*. Note that it preserves H and Θ_τ .

Let us set $H_{e,\tau} = A_{e,\tau} \cap H$. For every number $r \in (0, 1) \cap (\mathbb{Z} + \mathbb{Z}\theta)$ let us set

$$\mathcal{H}_{r,\tau} = \cup_{e:\text{tr}(e)=r} H_{e,\tau},$$

where the union is taken over the set of all idempotents e with $\text{tr}(e) = r$. Since $A_{1-e,\tau} = A_{e,\tau}$ we have $\mathcal{H}_{1-r,\tau} = \mathcal{H}_{r,\tau}$.

Theorem 2.17. *Assume that θ is irrational. Then*

$$\Theta_\tau = \cup_{r \in (0, 1/2) \cap (\mathbb{Z} + \mathbb{Z}\theta)} \mathcal{H}_{r,\tau}. \quad (2.6)$$

This decomposition is irreducible, i.e., none of the subsets is contained in the union of the rest. Also if $e \in A_\theta$ is any idempotent with $\text{tr}(e) = r$ then we have a map

$$\psi_e : (A_\theta^*)_0 \times H_{e,\tau} \rightarrow : (x, a) \mapsto x^{-1}ax + L(x) \quad (2.7)$$

such that $\mathcal{H}_{r,\tau}$ is the image of ψ_e . The codimension of the differential of ψ_e at a point (x, a) is equal to $\dim \text{Hom}_{\mathcal{C}}(E_1, E_2) + \dim \text{Hom}_{\mathcal{C}}(E_2, E_1)$, where $\mathcal{C}$ is the category of holomorphic bundles on $T_{\theta,\tau}$, E_1 and E_2 are the summands of the holomorphic decomposition $(A_\theta, \delta_\tau + \psi(x, a)) = E_1 \oplus E_2$ corresponding to the idempotent $x^{-1}ex$.

Proof. From the main theorem in [4] we know that a holomorphic bundle E on a noncommutative torus has non-scalar endomorphisms then it is either decomposable or its rank is a non-primitive element of $\mathbb{Z} + \mathbb{Z}\theta$. Since for a holomorphic bundle $E_a = (A_\theta, \delta_\tau + a)$ the second alternative is impossible, we conclude that $a \in \Theta_\tau$ iff E_a is decomposable. This implies (2.6). Since ψ_e is $(A_\theta^*)_0$ -equivariant with respect to the twisted action on H and the action by left multiplication on $(A_\theta^*)_0$, it suffices to consider the differential of ψ_e at a point $(1, a_0)$, where $a_0 \in H_e$. This differential maps $(x, a) \in A_\theta \oplus H \cap (eAe + (1-e)A(1-e))$ to $\delta_\tau(x) + [a_0, x] + a = d_{a_0}(x) + a$. But by Lemma 2.4

$$\text{coker}(d_{a_0} : A_\theta \rightarrow A_\theta) \simeq \text{Ext}_{\mathcal{C}}^1(E, E),$$

where $E = (A_\theta, \delta_\tau + a_0)$. Furthermore, it is easy to see that the map d_{a_0} is compatible with the decomposition

$$A_\theta = eA_\theta e + (1-e)A_\theta(1-e) \oplus eA_\theta(1-e) \oplus (1-e)A_\theta e$$

and that the cokernel of d_{a_0} acting on each of these four pieces is identified with $\text{Ext}^1(E_i, E_j)$ where $E = E_1 \oplus E_2$ is the holomorphic decomposition induced by e . The assertion immediately follows from this. $\square$

Let r_0 be the unique number of the form $\pm\theta + n$ with $n \in \mathbb{Z}$ such that $0 < r_0 < 1/2$.

Proposition 2.18. *One has*

- (i) $\Theta_\tau^{(1)} = \Theta_\tau$;
- (ii) $\Theta_\tau^{(2)} = \cup_{r \in (0, 1/2) \cap (\mathbb{Z} + \mathbb{Z}\theta), r \neq r_0} \mathcal{H}_{r,\tau}$;
- (iii) *the natural map*

$$\sqcup_{e:\text{tr}(e)=r_0} H_{e,\tau} \setminus \Theta_\tau^{(2)} \rightarrow \Theta_\tau \setminus \Theta_\tau^{(2)}$$

is an affine line bundle.

Proof. (i) follows from the fact that if E_a is decomposable then $\dim \text{End}_{\mathcal{C}}(E_a) > 2$. Moreover, the only case when $\dim \text{End}_{\mathcal{C}}(E_a) = 3$ is the following: $E_a \simeq E_1 \oplus E_2$ in the holomorphic category, where E_1 and E_2 are simple, $\text{Hom}_{\mathcal{C}}(E_2, E_1) = 0$ and $\text{Hom}_{\mathcal{C}}(E_1, E_2)$ is one-dimensional. Let $\text{rk } E_1 = m\theta + n$. Then $\text{rk } E_2 = -m\theta + (1 - n)$ and hence

$$\dim \text{Hom}_{\mathcal{C}}(E_1, E_2) - \dim \text{Hom}_{\mathcal{C}}(E_2, E_1) = -mn - m(1 - n) = -m$$

(see Corollary 2.9 of [4]). This implies that $m = -1$, so either $\text{rk } E_1 = r_0$ or $\text{rk } E_2 = r_0$ which proves (ii). It remains to note that the set of all possible holomorphic decomposition of E_a into E_1 and E_2 is a principal homogeneous space for $\text{Hom}_{\mathcal{C}}(E_1, E_2)$ which leads to (iii). $\square$

The part of Theorem 2.17 concerning tangent maps should imply that $\mathcal{H}_{r,\tau}$ has (in an appropriate sense) codimension ≥ 2 for $r \neq r_0$ (where $0 < r < 1/2$) while $\mathcal{H}_{r_0,\tau}$ is irreducible of codimension 1. If we were in the finite-dimensional situation we would immediately deduce from this that the hypersurface Θ_τ is irreducible. Our lack of knowledge does not allow us to state this precisely. However, we conjecture that the above statements hold when we intersect these loci with generic finite-dimensional subspaces in A_θ .

REFERENCES

- [1] K. R. Davidson, *C*-Algebras by Example*, Fields Institute Monographs, 6, AMS, Providence, RI, 1996.
- [2] K. Furutani, *On the Quillen determinant*, preprint math.DG/0309127.
- [3] A. Polishchuk, *Noncommutative two-tori with real multiplication as noncommutative projective varieties*, J. Geometry and Physics 50 (2004), 162–187.
- [4] A. Polishchuk, *Classification of holomorphic bundles on noncommutative two-tori*, preprint math.QA/0308136.
- [5] A. Polishchuk, *Quasicoherent sheaves on complex noncommutative two-tori*, in preparation.
- [6] A. Polishchuk, A. Schwarz, *Categories of holomorphic bundles on noncommutative two-tori*, Comm. Math. Phys. 236 (2003), 135–159.
- [7] D. Quillen, *Determinants of Cauchy-Riemann operators on Riemann surfaces*, Funct. Anal. and its Appl. 19 (1985), 37–41.
- [8] M. Rieffel, *The cancellation theorem for projective modules over irrational rotation C*-algebras*, Proc. London Math. Soc. (3) 47 (1983), 285–302.
- [9] M. Rieffel, *Noncommutative tori—a case study of noncommutative differentiable manifolds*, in *Geometric and topological invariants of elliptic operators (Brunswick, ME, 1988)*, 191–211, Contemp. Math. 105, AMS, Providence, RI, 1990.
- [10] L. Schweitzer, *Spectral invariance of dense subalgebras of operator algebras*, Int. Journal of Math. 4 (1993), 289–317.